

## **Patient Group Direction for**

# **Administration of Triamcinolone Acetonide 40mg/mL Injection in Peripheral Musculoskeletal Disorders by Podiatrists & Physiotherapists**

**Valid from: 1st October 2023  
Review date: 31<sup>st</sup> August 2025  
Expiry date: 1st October 2025**

**Version number: 9**

THIS PATIENT GROUP DIRECTION HAS BEEN AGREED BY THE  
FOLLOWING ORGANISATIONS:

**East Coast Community Healthcare  
and  
Norfolk and Waveney ICB**

**YOU MUST BE AUTHORISED BY NAME, UNDER THE CURRENT VERSION  
OF THIS PGD BEFORE WORKING UNDER IT**

**CLINICAL DETAILS**

<p><b>Indication</b></p>	<p>Treatment of adults with local inflammation of peripheral joints and soft tissues that would benefit from the use of local glucocorticoid. Suppression of local, peripheral inflammation and pain caused by soft tissue, joint and skin conditions.</p> <p>Frequently, Triamcinolone acetate will be administered alongside a local anaesthetic. Where this is the case, physiotherapist must also refer to and adhere to the Patient Group Direction (PGD) for that local anaesthetic.</p> <p>All practitioners working under this PGD must administer Triamcinolone and local anaesthetic separately and without mixing.</p> <p>Note: A PGD cannot be used for unlicensed medicines. Mixing of two or more licensed medicines constitutes the manufacture of a new unlicensed product. (Chartered Society of Physiotherapy (CSP) 2016)</p>
<p><b>Inclusion criteria</b></p>	<p>Patients aged 18 years and over that have given valid informed consent with benign peripheral joint or soft tissue pain, within the foot and/or ankle, including:</p> <ul style="list-style-type: none"> <li>• Rheumatoid arthritis or osteoarthritis with an inflammatory component in any joint</li> <li>• Bursitis</li> <li>• Capsulitis</li> <li>• Isolated soft tissue pathology of musculoskeletal origin</li> <li>• Entrapment neuropathy</li> <li>• Impingement syndromes</li> <li>• Ganglia</li> <li>• Ligamentous injury/lesions</li> <li>• Tendonitis/Tendinosis/Tendinopathy</li> <li>• Tenosynovitis</li> <li>• Enthesopathy</li> <li>• Myofascial Pain</li> <li>• Synovitis</li> </ul>

<p><b>Exclusion criteria</b></p>	<ul style="list-style-type: none"> <li>• Patients under the age of 18 years old</li> <li>• Pregnancy</li> <li>• Breast feeding</li> <li>• Severe / manic depression</li> <li>• Stomach ulcer</li> <li>• Presence of active local or systemic infection, including Tuberculosis (within the last 10 years)</li> <li>• Presence of symptoms suggestive of COVID-19: Cough, Temperature <math>\geq</math> 37.8, Loss of smell or taste, Flu-like symptoms</li> <li>• Adjacent osteomyelitis</li> <li>• Hypersensitivity / allergic reactions to corticosteroids or any of product components.</li> <li>• Recent trauma (six weeks) to the area intended for injection including hemarthrosis</li> <li>• Wound near to the injection site and fractures</li> <li>• Not into a Prosthetic joint itself or another site if joint replacement</li> <li>• Surgery performed in the area to be injected within the last 12 months</li> <li>• Planned surgical procedures (time scales depend on proposed injection and surgical procedure)</li> <li>• Unstable joints i.e. ligamentous insufficiency</li> <li>• Avascular area e.g., Achilles tendon</li> <li>• Presence of steroid arthropathy, poor skin condition over the area.</li> <li>• Immunosuppressed patients</li> <li>• Reluctant / non consenting patients</li> <li>• Unstable blood pressure</li> <li>• Unstable heart failure</li> <li>• Peripheral vascular disease at site to be injected</li> <li>• Severe or unstable heart condition</li> <li>• Sickle Cell disease</li> <li>• Metastatic carcinoma</li> <li>• Patients taking medication with potentially significant interactions with Triamcinolone:</li> </ul> <p><u>These include:</u></p> <p>Antibacterial drugs (e.g., erythromycin, clarithromycin), antifungals (e.g., ketoconazole), antivirals (e.g., HIV treatments), ciclosporin, and within 4 weeks of the administration of a live vaccine (e.g., yellow fever, MMR, BCG)</p> <p>➔ This list is not exhaustive - refer to the current British National Formulary (BNF) and/ or Summary Product Characteristics (SPC) for full list of drug interactions.</p>
<p><b>Precautions</b></p>	<p>Triamcinolone should be administered with caution in patients with the following conditions. Adverse effects are usually associated with systemic use at significant doses.</p>

Patients should be informed that their condition can be adversely affected by steroids, however, localised injections at the doses in this PGDs are safe and well tolerated. Refer to SmPC (Home - electronic medicines compendium (emc)) and discuss with a doctor or prescriber if you have concerns.

- Adrenal suppression / Cushing syndrome
- Poorly controlled/unstable Diabetes Mellitus
- Broken skin, eczema, history of infection in the area or adjacent to the area to be injected
- Glaucoma/Corneal Perforation
- Recent Implants e.g. mitral valve replacement, pacemaker
- Osteoporosis
- Liver failure or cirrhosis
- Renal impairment
- Myasthenia Gravis
- Epilepsy
- Anxious patients with signs of vagal over activity / needle phobia
- If COVID-19 vaccination is imminent, it may be appropriate to delay a non-essential steroid injection, as part of a shared decision, so that the response to the vaccine is more effective
- Bleeding disorders or anticoagulant therapy (Note: warfarin is in the exclusion criteria above)
- Concurrent steroid use (standard doses of inhaled corticosteroid are not significant)
- Ocular Herpes Simplex
- Peripheral vascular disease

**Anticoagulants:**

Warfarin (or any other Vitamin K antagonist):

- Patients must be on a stable dose with an INR result between 2 and 3 in the 7 days prior to treatment.
- It is advisable to check the INR again 3-4 days after the injection as corticosteroids may increase or decrease anticoagulant action

Direct Oral Anti-Coagulants (DOAC):

- Rivaroxaban, Apixaban, Edoxaban and Dabigatran
- Patients should not have taken their DOAC in the 12 hours preceding their injection
- The DOAC can be resumed at the previous dosing schedule when their next dose is due

Enzyme inducing medications including antiepileptic drugs (e.g., carbamazepine; phenobarbital); Antiviral drugs (ritonavir; efavirenz and nevirapine)

Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL  
Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0

	<p>cytotoxic drugs (e.g., methotrexate)</p> <p>➔ Always refer to accompanying product information leaflet</p> <p>➔ Therapy with triamcinolone does not obviate the need for the conventional measures usually employed. Although this method of treatment will ameliorate symptoms, it is in no sense a cure and the steroid has no effect on the cause of the inflammation.</p>
<b>Management of excluded patients</b>	<p>Consider alternative treatment. Document in patient physiotherapy record. Inform patient's GP and referrer (if different) in discharge summary or earlier if appropriate.</p> <p>If no alternative treatment within physiotherapy available, refer to General Practitioner with a Special Interest (GPwSI), Orthopaedic Consultant, Pain Clinic, Back Pain Service, other provider or back to GP depending clinical relevance and patient choice.</p>
<b>Action for patients not wishing to receive care under this PGD</b>	<p>Consider alternative treatment. Document in patient physiotherapy record. Inform patient's GP and referrer (if different) in discharge summary or earlier if appropriate.</p> <p>If no alternative treatment within physiotherapy available, refer to GPwSI, Orthopaedic Consultant, Pain Clinic, Back Pain Service, other provider or back to GP depending clinical relevance and patient choice.</p>
<b>DESCRIPTION OF TREATMENT</b>	
<b>Name of medicine</b>	Triamcinolone Acetonide 40mg/mL (Kenalog®) intra-articular/intramuscular injection
<b>Formulation and route</b>	<p>Using aseptic technique:</p> <ul style="list-style-type: none"> <li>• Intra-articular injection</li> <li>• Peri-articular</li> <li>• Soft Tissue Injection</li> </ul> <p><b>DO NOT ADMINISTER VIA THE INTRATHECAL OR INTRAVENOUS ROUTES</b></p>
<b>Strength</b>	40mg/ml
<b>Dosage</b>	<p>Single injection</p> <p>The following are dose guides for adults dependent on the specific disease entity being treated, the size of the joint and the severity of the pain:</p> <p>Recommended doses for intra-articular injection:</p> <ul style="list-style-type: none"> <li>• Up to 10 mg (0.25 ml) for a small joint</li> <li>• Up to 40 mg (1 ml) for a medium to larger joint</li> </ul>

Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0

	<p>Recommended dose for periarticular, intrabursal and tendon sheath injections:</p> <ul style="list-style-type: none"> <li>• 4 – 20 mg (0.1 – 0.50 ml)</li> </ul> <p>Specific recommendations for dosages can be found within Saunders and Longworth (2016) and the summary of Product Characteristics</p>
<b>Repeated dose instructions</b>	<p>Maximum of 3 injections per site (joint) per year</p> <p>Repeat injections are used as infrequently as possible in accordance with accepted clinical practice. Weight bearing structures being injected at no less than 4 monthly intervals.</p> <p>Referral for specialist opinion should be considered when the patient's condition does not respond positively to the injection.</p>
<b>Duration of treatment</b>	See sections above and below
<b>Quantity to administer</b>	<ul style="list-style-type: none"> <li>• The <b>maximum single dose</b> to be used per joint is 40 mg (1 ml)</li> <li>• The <b>total maximum dose</b> to be used across two or more joints per treatment episode is 80 mg of steroid (total volume 2 ml)</li> </ul> <p><b>Injections are not to be repeated if there is no benefit or change in condition.</b></p> <p>Referral for specialist opinion should be considered when the patient's condition does not respond positively to the injection.</p>
<b>Legal status</b>	Prescription only medicine (POM)
<b>Special Precautions</b>	<p>Always check patient's allergy status before treatment.</p> <p>Facilities for treating anaphylaxis must be available.</p>
<b>Adverse effects</b>	<p><b>Potential adverse drug reactions:</b></p> <ul style="list-style-type: none"> <li>• Transient flushing and dizziness</li> <li>• Infection</li> <li>• Anaphylaxis</li> <li>• Hyper or Hypopigmentation</li> <li>• Bleed / bruise</li> <li>• Subcutaneous atrophy</li> <li>• Post-injection flare</li> <li>• Irregular menstration</li> <li>• Severe joint destruction with necrosis of bone may occur if repeated intra-articular injections are given over a long period of time</li> </ul> <p><b>This list is not exhaustive. Refer to BNF and SPC via <a href="http://www.medicines.org.uk/emc">www.medicines.org.uk/emc</a> for complete list.</b></p> <p>Use the Yellow Card System via <a href="http://www.mhra.gov.uk/yellowcard">www.mhra.gov.uk/yellowcard</a> to report adverse drug reactions directly to the MHRA. Yellow Cards and guidance on its use are available at the back of the BNF.</p>

<b>Advice to be given</b>	<ul style="list-style-type: none"><li>• Assessment and preceding interventions will lead to a clinical decision that injection therapy may be of benefit to the patient. This clinical reasoning, alongside evidence-based practice, will be used to identify to the patient the suspected cause of their symptoms and why an injection may be of benefit as part of gaining valid informed consent.</li><li>• The risks and benefits associated with injection therapy including the possible adverse effects listed above will be discussed with the patient including what to watch for in relation to anaphylaxis and infection (see below)</li><li>• Advice regarding what to watch for in relation to anaphylaxis (itching acute skin changes, swelling, difficulty breathing, nausea, confusion), and infection (marked increase in pain accompanied by local swelling, further restriction of joint motion, fever, and malaise)</li><li>• That there is a potential risk of increased risk of and lack of clear evidence relating to COVID-19 and steroid injection</li><li>• Patient will be given a patient information leaflet (either Podiatry Department “Cortisone Advice Sheet”, or Physiotherapy Department “Information for Patients Considering Corticosteroid Injection Therapy”), to reinforce what has been discussed and/or product information leaflet. The leaflet will include details of local and systemic side effects.</li><li>• Patient will be given a patient information leaflet, “Corticosteroid Injection Therapy: COVID-19 Information Sheet”</li><li>• Valid informed written consent will be gained prior to an injection being performed and a copy of the consent form offered to the patient.</li><li>• The patient will be advised that they can stop the procedure and withdraw their consent at any point during the procedure.</li><li>• The patient will be advised about ‘relative rest’ and how to manage between injection and follow-up appointment, advice will depend on clinical condition and aims of treatment.</li><li>• Following the procedure, the patient will be advised to remain in department for 30 minutes following injection to assess and monitor any adverse reactions.</li><li>• The patient will be reminded of what to look for in relation to infection and advised to contact their GP for immediate assistance should infection occur.</li><li>• A follow up appointment will be arranged, and the patient will be given a contact telephone number and advised to contact the department should they have any queries or concerns before their follow up appointment.</li></ul>
---------------------------	--

Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL  
Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0

	<ul style="list-style-type: none"> <li>• Patients will be given the manufacturer’s patient information leaflet and advised about possible side-effects.</li> <li>• Patients should be advised to report any blurred vision or other visual disturbances (a rare retinal disorder associated with corticosteroid treatment).</li> <li>• Patients should be specifically warned to avoid over-use of joints in which symptomatic benefit has been obtained.</li> </ul>
<b>RECORDS AND FOLLOW UP</b>	
<b>Referral arrangements</b>	Refer to supervising doctor / facility or Accident & Emergency Department as appropriate
<b>Records to be kept</b>	<ul style="list-style-type: none"> <li>• Patient’s name, address, date of birth.</li> <li>• Contact details of GP (if registered).</li> <li>• Diagnosis or Working diagnosis</li> <li>• Consent given</li> <li>• Name of medication.</li> <li>• Dose and form (amount/ strength/ total volume).</li> <li>• Batch number and expiry date.</li> <li>• Site injected &amp; technique / approach</li> <li>• Advice given to patient (including side effects).</li> <li>• Signature/name of staff who administered or supplied the medication.</li> <li>• Copy of the signed consent form</li> <li>• Details of any adverse drug reaction and actions taken including documentation in the patient’s medical record.</li> <li>• Referral arrangements (including self-care).</li> <li>• Make entry into the patient’s records on SystemOne®, as appropriate.</li> <li>• A record of patients receiving injection therapy will be kept on system1 and can be identified by Business Intelligence, alongside an MSk Services audit document for review purposes</li> </ul>



<b>Follow up</b>	<ul style="list-style-type: none"><li>• A follow up appointment or telephone review will be booked with the patient at a time that is mutually acceptable for the patient and practitioner to assess the outcome of treatment.</li><li>• The patient will be advised on what to look for in relation to infection and advised to contact their GP for immediate assistance should infection occur.</li><li>• Timescales for the follow up appointment will be determined according to clinical relevance (usually two weeks post injection).</li><li>• The patient will be given a contact telephone number and advised to contact the department should they have any queries or concerns before their follow up appointment.</li><li>• Outcome of treatment will determine future management and the care pathway will be discussed with the patient as part of the valid informed consent process. Follow-up treatment may involve further injection therapy, referral to secondary care provider or back to GP. Patient choice and valid informed consent processes will be complied with at all times.</li><li>• Patient choice and valid informed consent processes will be complied with at all times</li><li>• The patient's medical practitioner will be informed of the injection procedure that includes date, drug name, dosage and site of administration in a timely manner</li></ul>
------------------	---

PROFESSIONAL REQUIREMENTS	
<b>STAFF CHARACTERISTICS</b>	<p>Qualifications required:</p> <ul style="list-style-type: none"> <li>Registered Podiatrist / Physiotherapist with a current Health Care Professions Council registration</li> <li>Experienced extended scope podiatrist in injection therapy OR any Podiatrist / Physiotherapist who has completed a corticosteroid injection therapy course accredited by Royal College of Podiatry / Chartered Society of Physiotherapists OR is working towards a recognised certificate in injection therapy (requiring at least 10 mentored injections) whilst under the supervision of a named mentor</li> </ul> <p>Additional requirements:</p> <ul style="list-style-type: none"> <li>Has undertaken appropriate training/education for working under patient group directions for the supply and administration of medicines.</li> <li>Understanding of pharmacology of drugs issued to patients and relevant medical condition.</li> <li><b>Immediate</b> access to Adrenaline/Epinephrine 1:1000 (1mg/mL) injection and trained in the treatment of anaphylaxis.</li> <li>Access to Resuscitation Council (UK) Emergency Treatment of anaphylactic reactions Guidelines for healthcare providers <a href="https://www.resus.org.uk/anaphylaxis/emergency-treatment-of-anaphylactic-reactions/">https://www.resus.org.uk/anaphylaxis/emergency-treatment-of-anaphylactic-reactions/</a></li> </ul> <p>Continued training requirements:</p> <ul style="list-style-type: none"> <li>Annual mandatory training on cardio-pulmonary resuscitation (CPR) and anaphylaxis.</li> <li>It is the responsibility of the individual to maintain competence and skills required to administer the medicine included in this PGD.</li> <li>A commitment to relevant CPD to include updating of relevant knowledge from current edition of British National Formulary (BNF) and awareness of any change to the recommendations for the medicine listed.</li> <li>It is the responsibility of the individual to keep up to date with continued professional development and to work within the limitations of individual scope of practice</li> </ul>

## REFERENCES / RESOURCES

### Summary of Product Characteristics Kenalog 40mg/ml injection

<https://www.medicines.org.uk/emc/product/6748/smpc> (Updated 8th April 2022 accessed 22/08/2023)

**Chartered Society of Physiotherapy** <https://www.csp.org.uk/professional-clinical/professional-guidance/medicines-prescribing-and-injections>

**Injection Techniques in Musculoskeletal Medicine:** A Practical Manual For Clinicians in Primary and Secondary Care. Saunders and Longworth, Churchill Livingstone Elsevier, 2016

**The British Society for Rheumatology:** Clinical guide during the COVID-19 pandemic for the management of patients with musculoskeletal and rheumatic conditions who are:  
-already taking corticosteroids, or-require initiation of oral/IV corticosteroids, or-require an intra-articular or intra-muscular corticosteroid injection 2020

[https://www.rheumatology.org.uk/Portals/0/Documents/COVID-19/MSK\\_rheumatology\\_corticosteroid\\_guidance.pdf](https://www.rheumatology.org.uk/Portals/0/Documents/COVID-19/MSK_rheumatology_corticosteroid_guidance.pdf)

**The British Society for Rheumatology: COVID-19:** Guidance For Rheumatologists 2021  
<https://www.rheumatology.org.uk/practice-quality/covid-19-guidance>

## PGD DEVELOPMENT

### Details of Patient Group Direction owner

Name: Mr Martin MacDonald  
Position: Extended Scope Practitioner and Team Lead MSk Triage Service  
Contact Email: martin.macdonald@ecchcic.nhs.uk  
Contact Address: East Coast Community Healthcare (ECCH)

### Patient Group Direction peer reviewed on behalf of ECCH by

Name	Position	Date
	Physio/ Podiatry development virtual group	
	Medicines Management Group	

**This patient group direction must be agreed to and signed by all health care professionals involved in its use. East Coast Community Health Care CIC should hold the original signed copy. The PGD must be easily accessible in the clinical setting**

### Organisation

East Coast Community Healthcare

Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL  
Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0

<b>Lead Doctor</b>	<p>Name: Dr Rupert Talboys</p> <p>Position: GP Associate</p> <p>Signature:</p> <p><i>wet signature</i></p> <p>Date:</p>
<b>Lead Pharmacist</b>	<p>Name: Ms Françoise Price</p> <p>Position: Head of Pharmacy and Medicines Optimisation</p> <p>Signature:</p> <p><i>wet signature</i></p> <p>Date:</p>
<b>Lead Practitioner of the professional group to supply medicines under this PGD</b>	<p>Name: Mr Martin MacDonald</p> <p><b>Position:</b> Extended Scope Practitioner and Team Lead MSk Triage Service</p> <p>Signature:</p> <p><i>wet signature</i></p> <p>Date:</p>
<p><b>This PGD has been agreed to be appropriate, necessary and an advantage to patient care by the East Coast Community Healthcare Medicines Management Group</b></p>	

## PGD AUTHORISATION

**This PGD is authorised by:**

**NHS NORFOLK AND WAVENEY ICB**

**Director of Nursing**

Name: Tricia D'Orsi

Signature: 

Date 18/10/2023

**Associate Director of  
Pharmacy and  
Medicines Optimisation  
(Chief Pharmacist)**

Name: Michael Dennis

Signature: 

Date: 20/10/2023

## INDIVIDUAL AUTHORISATION

**YOU MUST BE AUTHORISED BY NAME, UNDER THE CURRENT VERSION  
OF THIS PGD BEFORE WORKING UNDER IT**

BY SIGNING THIS PATIENT GROUP DIRECTION YOU ARE INDICATING THAT YOU AGREE TO  
ITS CONTENTS AND THAT YOU WILL WORK WITHIN IT

**PGDs DO NOT REMOVE INHERENT PROFESSIONAL OBLIGATIONS OR ACCOUNTABILITY**


**IT IS THE RESPONSIBILITY OF EACH PROFESSIONAL TO PRACTICE ONLY WITHIN THE  
BOUNDS OF THEIR OWN COMPETENCE**

**IF THIS IS AN UPDATED OR REPLACEMENT PGD ENSURE THAT ALL OLDER VERSIONS ARE  
WITHDRAWN FROM USE WITH IMMEDIATE EFFECT**

**IT IS YOUR REponsibility TO MAKE SURE YOU ARE USING THE CURRENT VERSION**

*NOTE TO AUTHORISING MANAGERS: AUTHORISED STAFF SHOULD BE PROVIDED WITH AN  
INDIVIDUAL COPY OF THE CLINICAL CONTENT OF THE PGD AND A PHOTOCOPY OF THE  
AUTHORISATION SHEET SHOWING THEIR AUTHORISATION. BY SIGNING BELOW YOU ARE  
CONFIRMING THAT YOU HAVE ASSESSED THE STAFF MEMBER AS COMPETENT TO WORK  
UNDER THIS PGD AND THAT THEY HAVE THE ORGANISATIONAL APPROVAL TO DO SO*

Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL  
Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0



Patient Group Direction for Administration of Triamcinolone Acetonide 40mg/mL  
Injection in Peripheral Musculoskeletal Disorders by Podiatrists and Physiotherapists  
Version: 9.0

<b>Name of Professional</b>	<b>Signature</b>	<b>Authorising Manager</b>	<b>Date</b>